



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/47, C12Q 1/68, C07K 14/705</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/35878</b> <b>(43) International Publication Date:</b> 2 October 1997 (02.10.97)
<b>(21) International Application Number:</b> PCT/NL97/00157 <b>(22) International Filing Date:</b> 27 March 1997 (27.03.97) <b>(30) Priority Data:</b> 96200855.3 28 March 1996 (28.03.96) EP <i>(34) Countries for which the regional or international application was filed:</i> NL et al. <b>(71) Applicants (for all designated States except US):</b> DALLAND B.V. [NL/NL]; Postbus 128, NL-5740 AC Beek en Donk (NL). PROVA B.V. [NL/NL]; Postbus 290, NL-5280 AG Boxtel (NL). STAMBOEK ZUID B.V. [NL/NL]; Postbus 86, NL-5268 ZH Helvoirt (NL). NOORD NEDERLANDS VARKENSSTAMBOEK B.V. [NL/NL]; Hoofdweg 2, NL-7382 BH Klarenbeek (NL). INSTITUUT VOOR DIER- HOUDERIJ EN DIERGEZONDHEID (ID-DLO) [NL/NL]; Edelhertweg 15, NL-8219 PH Lelystad (NL). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> GERBENS, Frans [NL/NL]; Botter 25-02, NL-8232 KP Lelystad (NL). <b>(74) Agent:</b> SMULDERS, Th., A., H., J.; Vereenigde Octrooibureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> THE PORCINE HEART FATTY ACID-BINDING PROTEIN ENCODING GENE AND METHODS TO IDENTIFY POLYMORPHISMS ASSOCIATED WITH BODY WEIGHT  <b>(57) Abstract</b>  The present invention provides a novel sequence of the pig H-FABP gene, as well as methods of using said gene and its products. Especially the invention provides methods for detecting different alleles of the pig H-FABP gene, which different alleles are associated with differences in the genotypic and/or phenotypic traits of the pigs having those alleles. Especially the invention provides methods for distinguishing between alleles resulting in different phenotypes, particularly using techniques involving selective amplification of pig H-FABP gene derived materials. These techniques are especially suitable for selecting animals to be used in breeding programmes. Breeding programmes employing such techniques are also disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

THE PORCINE HEART FATTY ACID-BINDING PROTEIN ENCODING GENE AND METHODS TO  
IDENTIFY POLYMORPHISMS ASSOCIATED WITH BODY WEIGHT

INTRODUCTION

The invention relates to the field of molecular biology as well as to the field of breeding methods for farm animals, in particular pigs. In particular the invention relates to the use of diagnostic methods derived from the field of molecular biology to be applied in breeding programmes that select animals on production traits that improve their breeding value.

By selecting animals on their breeding value calculated mainly from phenotypic measurements of production traits, breeding has greatly improved the genotype for production traits of livestock animals. Thus, traditionally, breeding programmes have selected for phenotypic characteristics of animals. However, more recently selection for genotypic characteristics that are associated with improved production traits have gained interest in the field. Selection for phenotypic characteristics entails mainly selection of the offspring or siblings or other relatives of the animals to be selected whereas selection of specific genotypic characteristics allows for earlier and specific detection of animals of interest.

Within methods that select on specific genotypic characteristics, one may distinguish between methods that detect genetic variation in genes or quantitative trait loci that are merely associated with production traits of animals and methods that detect genetic variation in functional genes that directly influence those production traits. One of the former methods is a marker assisted selection wherein polymorphisms in markers identified in a random manner are associated with production traits.

For instance, meat production is closely linked to embryonic muscle formation, and, consecutively, to the distribution of muscle cells and fat cells. Biologically, production is concentrated in defined tissues of the animal, e.g. muscle tissue for lean meat production. In breeding programmes for optimizing porcine lean meat production, various levels of selection pressure have been applied to different tissues (i.e. muscle, fat and bone). However, when selecting for lean meat, and thus the absence of fat, one may lose certain traits that are wanted after all, i.e. traits that are associated with taste and thus with the consumers' perception of the final product.

In pig breeding programs traditionally a lot of emphasis has been put on the fat reduction because of the consumers interest in lean meat. Fat reduction is surveyed as a decrease in backfat thickness and a large reduction has been achieved since the establishment of breeding programmes in pigs. However reduction of the backfat depot also results in less intramuscular fat (IMF). This last depot is the main fat depot in meat and is positively correlated with the taste and thus the acceptance of meat (Wood et al., 1988). To exclude the IMF depot from further reduction a marker for this trait is necessary because IMF is hardly measurable in living animals. Recently, it has been statistically shown that a single major gene for IMF deposition in pigs must be present (Janss et al., 1994), however, the sequence, location and mode of action of the putative gene were not disclosed. Here we present evidence of a muscle tissue specific candidate gene located on porcine chromosome 6 which is the heart fatty acid-binding protein (H-FABP) gene. Genetic variation in this gene is responsible for the variation in among others IMF% and other production traits of pigs.

Fatty acid binding proteins (FABP's) are small intracellular proteins involved in fatty acid transport from the membrane to the sites of  $\beta$  oxidation and/or triacylglycerol or phospholipid synthesis (Veerkamp and

Maatman, 1995). Furthermore, FABP's modulated the intracellular fatty acid concentration (Veerkamp et al., 1993). Fatty acid metabolism has historically been linked to insuline resistance (Randle, 1963), and therefore mutations in FABP genes may be associated with changes in cellular insulin resistance or dependency, fatty acid oxydation and fatty acid binding. FABP's are members of a family of intracellular lipid binding proteins comprising at least eight structurally distinct types originating from: adipocytes, brain, epidermal cells, heart, intestinal cells, ileal cells, liver and myelin cells.

The heart type FABP (H-FABP) is a 15 kDa protein highly expressed in several tissues with a high demand for fatty acids, such as cardiac and skeletal muscle and lactating mammary gland. H-FABP is identical to MDGI (mammary derived growth inhibitor) a protein which inhibits growth of tumor cells *in vitro* (Bohmer et al., 1987). Functionally, the H-FABP can induce cardiac myocyte hypertrophy *in vitro*, when added to the culture (Burton et al., 1994) and also promotes functional differentiation of mammary epithelial cells *in vitro* (Yang et al., 1994). However, no secretion of H-FABP has been detected so far. On the other hand native and overexpression of H-FABP in mammary epithelium of lactating mice (*in vivo*) does not correlate with functional differentiation markers of these cells (Binas et al., 1995).

The present invention provides among others an isolated or recombinant pig H-FABP gene specific nucleic acid molecule or pig H-FABP gene specific fragments thereof comprising or hybridising to the nucleotide sequence as shown in figure 1, or its complementary sequence or the RNA equivalents thereof.

The locus of this gene is on porcine chromosome 6. The pig H-FABP gene can be assigned functions in the regulation of intramuscular fat, thereby changing the ratio of fat deposited within the muscle versus fat deposited outside the muscles, i.e. in backfat depots. Since production and deposition of fat is energy consuming and takes away energy

for other purposes, such as muscle growth, the regulation of intramuscular fat is correlated to the regulation of growth, and thus body weight and average daily gain and feed efficiency. Also, H-FABP can regulate myocyte (and thus muscle) hypertrophy and thus also muscle regeneration. Since FABP's are involved in fatty acid transport they can influence fatty acid oxidation rates, the metabolism of fatty acid derivatives in the tissue and the fatty acid composition of cells and thus of meat. Furthermore, FABP's may regulate cellular insulin dependency. Also, in pregnant animals, fat storage has an impact on embryo survival, and regulation of H-FABP will influence birth rates and litter size. Since H-FABP regulates functional differentiation of mammary epithelial cells it may be involved in regulating the quantity and composition of the milk available, thus influencing the growth and survival of newborn animals. With the present invention, the genetic variation within the pig H-FABP gene with respect to variation in regulation of expression can now be revealed and analysed for association with above production traits and physiological characteristics.

The present invention further provides a method to generate via recombinant DNA techniques an animal, such as small laboratory animals or farm animals, i.e. a pig, with additional genetic material originating from the pig H-FABP gene. Such animals may then encode wanted alleles of this gene and constitutively or transiently express allelic proteins or fragments thereof that enhance the production or physiological characteristics of those animals.

The invention further provides methods to generate proteins or (poly)peptides comprising various allelic proteins or fragments thereof derived from the pig H-FABP gene. Such peptides, or antibodies specifically directed against such peptides, may be used to influence production traits in the live animal, but may also be used in cell-culture systems *in vitro*. Such (poly)peptides or proteins, or antibodies specifically directed against these, may also

be used in diagnostic test systems to select animals that express wanted forms of allelic proteins or fragments thereof encoded by the pig H-FABP gene.

The invention further provides methods localising, identifying or marking genes or alleles or quantitative trait loci, in particular those corresponding to the pig H-FABP gene, in samples, in particular biological samples, cells or tissues, such as but not limited to hair, skin or blood, of farm animals, in particular pigs, by allowing for specific amplification of genomic fragments of those genes or alleles or quantitative trait loci of pigs. Since marker assisted selection of animals is frequently based upon genetic variation that exists within functional genes that influence a production trait directly, i.e. genes such as the pig H-FABP that regulates fatty acid binding, one of the methods that the invention provides is a method that identifies or marks loci or genes and that can distinguish between characteristics of alleles of those genes which characteristics serve as markers in selection programmes for animals with specific versions of those genes that are directly linked with improved production traits.

The invention further provides a method wherein polymorphic restriction sites within functional genes and thus different alleles of those genes are identified by allowing for specific amplification of genomic fragments of those genes, in particular by allowing for specific amplification of fragments of the H-FABP gene. Amplification methods are well known in the art, the best known being PCR. A short description of the PCR used herein is given in the experimental part. Other primers, enzymes and conditions can of course be applied. After amplification a suitable method of identifying wanted alleles is a restriction endonuclease treatment. Suitable restriction enzymes for pig H-FABP alleles are *MspI*, *HaeIII* or *HinfI*, but others may also be used. By these methods large numbers of pigs can be rapidly genotyped for studies in which genotypic variation can be

associated with growth characteristics and other production traits of pigs.

However, there are many other methods identifying polymorphisms in alleles, both at the nucleic (DNA/RNA) level and at the product (protein) level. In particular at the protein level there are many possibilities using immunoassays, whereas at the nucleic acid levels there are many assays which all include some kind of hybridisation step of for instance primers or labelled nucleic acids. A very good possibility would be mismatch PCR. Primers to be used in the invention can be identified by the person skilled in the art, the sets given in the experimental part are for illustrative purposes only.

Furthermore, the methods according to the invention can be developed into diagnostic assays or kits by which selection of pigs with alleles of interest can be performed in routine screening protocols employed in breeding programmes. With such protocols better results of selection can be expected when genes responsible for regulation of commercially interesting body tissues can be rapidly identified and controlled.

In the specific case of the pig H-FABP gene, such testing protocols can be used to identify, select and breed farm animals, such as pigs, which have better production traits, such as IMF% or backfat thickness or average daily weight gain or feed efficiency, than the average animal in the population. Better production traits such as BW or daily weight gain will increase the production per year expressed as amount of meat per animal raised. A population of animals with a higher and less variable IMF% will result in a more homogenous product (meat) which is also better appreciated by putative customers because of a better taste. Furthermore, selection for higher IMF% may be possible while at the same time selection against fat deposition in other depots, such as backfat, can be performed.



## EXPERIMENTAL

The porcine H-FABP gene has been isolated, characterized and chromosomally localized. Polymorphisms in this gene have been identified. To test the association between bodyweight (BW) and percentage of intramuscular fat (IMF %), animals with different polymorphisms were selected, their bodyweight was measured and the amount of IMF after slaughter was measured.

10

## MATERIALS AND METHODS

## Isolation of H-FABP containing phage clones.

15 A porcine genomic DNA EMBL3/SP6/T7 library (Clontech Laboratories Inc. Palo Alto, CA) was screened using the plaque hybridization method (Sambrook et al., 1989). The human H-FABP cDNA cloned in the pSP65 vector (Peeters et al., 1991) and labeled with  $^{32}\text{P}$ -dCTP by nick translation  
20 (Sambrook et al., 1989) was used as a probe. Briefly, 500.000 plaques were transferred to replica nitrocellulose filters and incubated in denaturation buffer (1.5 M NaCl, 0.5 M NaOH) for 2 minutes, neutralisation buffer (1.5 M NaCl, 0.5 M Tris-HCl pH 8.0) for 5 minutes and fixation  
25 buffer (0.2 M Tris-HCl pH 7.5, 2X SSC(0.3 M NaCl, 0.03 M Sodium citrate)) for 30 s. The filters were air dried and the DNA was irreversibly bound by baking the filters at 80°C for 2 h.

The filters were prehybridized (6X SSC, 0.5% (w/v) SDS, 30 5X Denhardt's and 100ug/ml NaOH treated salmon sperm DNA) for two hours at 67°C and hybridized in identical buffer with the addition of the radioactively probe at 67°C overnight. The filters were washed four times with 2X SSC, 0.1% (w/v) SDS for 30 min at room temperature. Twenty  
35 plaques that showed positive signals on both replica filters were isolated and each subjected to two additional rounds of low density plaque purification.

DNA of these clones was isolated using the plate lysate method (Sambrook et al., 1989).

### Polymerase chain reactions.

5

PCR amplifications were performed on 1  $\mu$ l of a 1:1000 dilution of phage DNA preparations or 50 ngr of genomic DNA in 50  $\mu$ l containing 0.2 U Super Tth polymerase (SphaeroQ, Leiden, NL) in 10 mM Tris-HCl pH 9.0, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.1% (w/v) gelatin, 1% Triton X-100, 0.5  $\mu$ M of each primer (Pharmacia Biotechnologies, Uppsala, Sweden) and 0.2  $\mu$ M of each dNTP (Boehringer Mannheim, Mannheim, Germany). After 3 min of denaturation at 94°C, 33 cycles of amplification were carried out: 94°C for 1 min, the indicated annealing temperature (Tables 1 and 2) for 1 min and 72°C for the time considering the length of the expected fragment (approximately 1 min for every kb).

The primer sequences as used for the poly-T microsatellite amplification are

[Hex]-5'TCTGGGCTTCAACTTACTCTG3' and  
5'CTAGCGCTTCAGCTCTGATTG3'

PCR was performed as described before with the following cycling conditions, 20 s at 94°C, 40 s at 57° and 1 s at 72°C followed by a final extension step for 10 min at 72°C, PCR products were analysed on a 8% polyacryl amide gel in an ABI373A (Perkin Elmer, Foster City, CA, USA) using Genescan 1.0/Genotyper 1.1.1.2 software (Perkin Elmer). Allele sizes were estimated by comparison with a commercial Tamra-labeled marker (Perkin Elmer).

### DNA sequence analysis

PCR#1 (Table 1) was performed on DNA of all the phage clones to detect porcine H-FABP intron 1 specific fragments. Two of the three phage clones containing the H-FABP gene were used to subclone the various SacI and KpnI (Boehringer

Mannheim, Mannheim, Germany) restriction digestion fragments of the gene region in pBS. Unfortunately, neither of these phage clones contained exon 4 and the 3' untranslated region as detected by restriction analysis.

5       The intron 3/exon 4 splice junction was cloned as PCR#2 (Table 1) product using porcine (Great Yorkshire) genomic DNA as a template. The products of two independent PCR reactions were cloned to identify errors by the Super Tth polymerase upon analysis.

10       The 3' untranslated region was isolated using the 5'/3' RACE-PCR kit (Boehringer Mannheim, Mannheim, Germany) with porcine (Meishan) muscle cDNA as the template and porcine H-FABP exon 1 or 3 (Table 1) specific primers in combination with the provided poly-A primer.

15       PCR products were cloned in the pT7Blue vector (Novagen Inc., Madison, U.S.A.).

20       All H-FABP (sub)clones were transformed and the recombinant plasmid DNA was isolated and purified with the Wizard Maxiprep kit (Promega, Madison, WI, U.S.A.). The nucleotide sequence was determined by dideoxy sequencing, partially by cycle sequencing (Perkin Elmer,) or autoread sequencing (Pharmacia Biotechnologies, Uppsala, Sweden) and the analysis was performed on a ABI 373 (Applied Biosystems) or ALF DNA sequenator (Pharmacia Biotechnologies, Uppsala, Sweden) respectively.

#### RFLP screening

30       Porcine genomic DNA was isolated as described (Sambrook et al., 1989) from EDTA treated blood stored at -80°C. One hundred ng of genomic DNA was used for PCR amplification in 50 µl reaction as described before. The primer sequences and its corresponding product size and annealing temperature for each combination are given in table 2. Fifteen µl of the PCR reaction was used for restriction digestion with 2 units of *HaeIII*, *HinfI* or *MspI* (Boehringer Mannheim) in a total volume of 20 µl. For *HaeIII* and *HinfI* the recommended buffer

conditions were additionally used whereas *MspI* was added directly to the PCR buffer. Restriction digestion fragments were loaded on a 2% (*MspI*) or 3% (*HaeIII* and *HinfI*) agarose (Sigma, St Louis, MO, U.S.A.) gel and after electrophoresis the RFLP patterns were scored by two persons, independently.

### Chromosomal localisation

Two independently established pig/rodent somatic cell hybrid panels (Panel A: Rettenberger et al., 1994a, 1994b, 1994c, 1995a and Panel B: Zijlstra et al., 1994a, 1994b, 1994c, in prep.) were used to assign the H-FABP gene to a specific chromosome by PCR.

DNA from each cell hybrid containing porcine chromosomes in various combinations was used in PCR#3 (Table 1) which unambiguously amplified porcine H-FABP intron 3 sequences. The obtained data was statistically evaluated according to Chevalet and Corpet (1986) in comparison with the cytogenetically and/or reference loci data of both panels.

### Analysis of bodyweight and intramuscular fat in relation to polymorphisms in H-FABP genotypes.

One hundred Duroc pigs were selected on the basis of the amount of IMF in their slaughtered relatives. The animals came from the test farms Someren (N=50) and Herpen (N=50). Blood samples were used to isolate DNA which was used to determine the genotypes of the three polymorphisms of each animal. Of all animals BW was measured. Animals which were not used in the regular breeding program were slaughtered and analyzed for the amount of IMF. Only 45 animals were slaughtered. For three of these animals it was not possible to determine the genotype of one or more polymorphisms. In total 42 animals were used in the analyses.

An additional experiment was performed to further study the production traits in relation to genetic variation in the H-FABP gene. Therefore thirteen boars and seventy-two dams were selected for this investigation from two Duroc populations housed at separate test stations. Selection was based on heterozygosity for each H-FABP PCR-RFLP. Progeny was housed in groups and fattened with ad libitum food access until slaughterweight (110 kg). Performance traits recorded for each pig were live weight at 180 days of age (BW), backfat thickness (BFT) and for each dam the number of piglets produced alive in first (FPP) or second parity (SPP), respectively. At slaughter, meat quality traits i.e. cooking loss, drip loss, intramuscular fat percentage, minolta colour, pH and shear force were measured in a subset of the slaughtered animals. Blood or hairroots were collected from each animal to isolate genomic DNA for H-FABP PCR-RFLP genotyping. The final dataset comprises information from in total 2345 pigs including pedigree. For 823 pigs H-FABP genotype information is available for at least one of the PCR-RFLPS.

Analysis was done with the statistical program Statistical Analyses System (SAS, 1990) and with the program Prediction and Estimation (PEST, Groeneveld, 1990). The latter program uses family information to estimate Best Linear Unbiased Predictions (BLUP) for the influence of, in this case different genotypes on the amount of IMF and BW. The mean amount of IMF% of the Duroc population is 3.20, its standard deviation is 0.84 (Hovenier, 1992). PEST was also used in combination with the GeneProb program (Kerr and Kinghorn, 1996) that estimates missing genotypes for animals based on genetical and or phenotypical information for the trait of interest. Thus, the effect of the H-FABP genotypes on performance, and meat quality traits was studied with the following model:

Trait= int + test station\*test year\*test month + sexe +  
litter + P(XX) + P(Xx) + P(xx) + individual + covariable +  
residual effect

5 where

P(XX), (Xx), (xx) are the estimated chance for each genotype  
for each animal.

BW was standarized to weight at 180 days (STD-BW)

BFT was standarized to a weight of 110 kg (STD-BFT)

10 For IMF age was included as a covariable and analysis was  
also performed with STD-BFT and STD-BW as covariables.  
Heritability ( $h^2$ ) estimates for each trait in this Duroc  
population were assumed to be similar to the estimates  
described by Hovenier et al. (1992) for the Duroc breed.

15

## RESULTS

### H-FABP gene sequence determination and analysis

20 Twenty H-FABP positive phage clones were identified and  
the corresponding DNA isolated and examined for the presence  
of the H-FABP gene. Using PCR#1 (Table 1) three phage clones  
appeared to contain intron 1. The rest of the phage clones  
contained H-FABP pseudogene-like sequences because of the  
25 absence of intron 1 and 3 in the amplification product of  
PCR#1 and #2 (Table 1) respectively. Sequence analysis of  
these pseudogene specific amplification products showed  
various nucleotide substitutions in comparison with the H-  
FABP gene coding sequences. Furthermore a 27 bp internal  
30 duplication was detected in the PCR#3 amplification product  
of one H-FABP pseudogene containing phage clone (data not  
shown). However this particular pseudogene specific PCR#3  
fragment was not detected in the main pig breeds of our  
panel.

35 Further PCR analysis of the three H-FABP gene  
containing phage clones revealed that neither contained the  
exon 4 and 3' untranslated region of the gene. Using PCR#2

on porcine genomic DNA a 1500 bp fragment was amplified and cloned for intron 3 and exon 4 sequence analysis. The 3' untranslated region (3'UTR) was amplified on porcine whole muscle cDNA, cloned and sequenced.

5       The coding sequences with the flanking intronic sequences and also 1600 bp of the 5' upstream region were determined (Fig.1). The exon-intron splice junctions were located in comparison with the porcine H-FABP cDNA and the murine H-FABP/MDGI (Treuner et al., 1994) gene sequence. A  
10       potential TATA-box was located 92 bp upstream the ATG start codon and in the (3'UTR) a consensus poly-A signal sequence was identified (see Fig.1). The coding sequences showed 92%, 91%, 87% and 85% identity to the bovine, human, mouse and  
15       rat H-FABP sequences at the nucleotide level and the deduced amino acid sequence were 92%, 90%, 87% and 86% identical, respectively (Billich et al., 1988; Peeters et al., 1991; Binas et al., 1992; Claffey et al., 1987).

#### Detection of genetic variation

20

A panel comprising genomic DNA of 7 pig breeds each represented by unrelated animals (see table 3) was used to detect genetic variation in the 5' upstream region, intron 2 and intron 3 of the porcine H-FABP gene. Therefore, part of  
25       the 5' upstream region was amplified on DNA of this panel using PCR (Table 2) and digested with the restriction enzyme *HinfI*. The *HinfI* digestion showed two alleles a single fragment of 256 bp (allele h) or two fragments of 197 and 59 bp (allele H). Similarly intron 3 (PCR#3, Table 1) and intron  
30       2 (Table 2) were tested for genetic variation with the enzymes *MspI* and *HaeIII* respectively, and both showed genetic variation in intron 2. *HaeIII* showed one fragment of 850 bp (allele D) and/or fragments of 400 and 450 bp (allele d). Accurate size determination revealed that these three  
35       fragments were 684 bp, 278 bp and 406 bp. *MspI* showed a fragment of 850 bp (allele a) and/or fragments of 750 and 50 bp (A). Accurate size determination revealed that these

fragments were 814 bp, 703 bp and 111 bp. Both sites of genetic variation are approximately 300 (285) bp apart.

Sequence analysis of the porcine H-FABP gene sequence (Fig. 1) revealed a 25 thymidine-nucleotide (poly-T  
5 microsatellite) stretch in the first intron. To investigate genetic variation in this poly-T stretch this region was amplified by PCR. In the Duroc pigs at least 3 alleles (H1, 215-bp, H2:220-bp and H3:221-bp) were detected. Obviously, these alleles showed a complete linkage with H-FABP PCR-RFLP  
10 alleles which are located within a 1,5 kb region.

Table 3 represents the allele frequencies of the different PCR-RFLPs in the different pig breeds tested.

The mendelian inheritance pattern of the three PCR-RFLPs was analysed in a porcine family comprising 3  
15 generations of a Great Yorkshire breed. The genotypes of the individual pigs show consistent patterns of inheritance in this family.

#### **Chromosomal localisation**

20

The porcine H-FABP gene was chromosomally localized using a porcine H-FABP gene intron 3 specific PCR which amplified no rodent homologous. Amplification on DNA of two independently established pig/rodent cell hybrid panels and  
25 comparison with the cytogenetically (panel A and B) and reference loci data (panel A) revealed a single significant association of the H-FABP gene with chromosome 6 (Table 4) for both cell hybrid panels.

#### **Analyses of bodyweight and intramuscular fat in relation to polymorphisms in H-FABP genotypes.**

Table 5 shows the result of mean values and their standard deviations of IMF and bodyweight for different  
35 fixed effects which were taken into account in this analyses.



### Statistical analyses using SAS

The data was analyzed using the General Linear Models procedure (GLM) from SAS (Sas, 1990). A large model (lrg) with IMF as dependant variable contained the (fixed) effects test farm, sexe, *MspI*, *HaeIII* and *HinfI*. Also models containing the different effects separately (ind) and one containing the combined genotype of these three polymorphisms were analyzed. Table 6 shows the significance values for the different effects for the models analyzed. Table 9 shows the effects of the three polymorphisms on intramuscular fat%.

### Statistical analyses using PEST

The PEST program was used to be able to use family information in the analyses of the different fixed effects. The used model contained the same fixed effect as the model with SAS but also contains a random animal effect. Also, a pedigree file was used containing family relations up to two generations back. Table 7, 10, 11, 12 and 13 show predicted values for the different fixed effects and their standard errors.

### 25 Hypothesis testing

The different values and their standard errors, found for the fixed effects where used in a Chi-squared hypothesis test. Critical values were calculated for a 90% and 95% two-sided confidence limit. Calculation was performed using formula 1 and 2. The  $\mu_1$ ,  $\mu_2$ , s.e.1 and s.e.2 are taken from PEST output.

The value  $t_{1-1/2\alpha}$  is taken from a confidence table and has a value of 1.96 for a 95% confidence interval.

35

$$\begin{aligned} y_1 - y_2 &= \mu_1 - \mu_2 \pm t_{1-1/2\alpha} \sqrt{(s.e.2)} \\ s.e.2 &= (n_1 \cdot s.e.1^2 + n_2 \cdot s.e.2^2) / (n_1 + n_2 - 2) \end{aligned}$$

$t_{1-1/2\tau}$  = 1.96 (95% confidence limits)  
 $\mu_1 - \mu_2$  = difference between two genotypes of one polymorphism

5 Table 8 shows the difference between different genotypes of each polymorphism and its 90% and 95% critical values.

The differences between the values for the different genotypes on IMF are not significant (90%). Only the  
10 difference between the homozygote combination (aaddHH) and the heterozygote combination (AaDdHh) showed a significant difference (90%) but was not significant with the 95% two sided confidence limits.

Bodyweight is significantly different for the different  
15 genotypes in all three polymorphisms.

When the different genotype combinations are analysed for bodyweight the heterozygote combination (AaDdHh) did not differ significantly from both homozygote genotypes. Both  
homozygote genotypes (AADDhh-aaddHH) show a significant  
20 difference of 9.11 kg ( $P < 0.05$ ).

## Conclusions

The genotypes of the three polymorphisms tested (*MspI*,  
25 *HaeIII* and *HinfI*) show a significant (95%) difference in bodyweight (BW). All three polymorphisms can be used in selection for bodyweight. The genotypes of the three polymorphisms show a distinct, albeit non-significant difference in IMF percentage. If there is a difference  
30 between different genotypes of 0.20, 50 animals of the least frequent genotype (AADDhh) would be needed to reach a significant (95%) difference of 0.2. Tables 9, 10 and 11 show that when more animals are tested, statistically significant differences among the three polymorphisms can  
35 indeed be found, for instance for IMF, backfat thickness and BW.

Also, tables 12 and 13 show that the effect on IMF, as measured by RFLP testing, can still be found when the effects are corrected for backfat thickness and/or growth.

### Brief description of the drawings

Figure 1. The porcine H-FABP gene sequence including  
5 1632 bp of the 5'upstream region and 200 bp of the 3'  
untranslated region. Exons are represented by bold capital  
letters and the deduced amino acid sequence is shown  
directly beneath it. Standard one letter amino acid symbols  
are used. The putative TATA-box, the polyadenylation signal  
10 in the 3'UTR and the 13 nucleotide element are underlined.  
The size of the nondepicted intron sequences is shown  
between arrowheads. The polymorphic *HaeIII* (GGCC), *HinfI*  
(GATTC), *MspI* (CAGG) sites and the polymorphic  
microsatellite sequence (poly-T) are depicted bold and  
15 underlined.

## REFERENCES

- Archibald, A.L., Haley, C.S., Brown, J.F., et al. (1995) The PIGMaP consortium linkage map of the pig (*Sus scrofa*). *Mamm. Genome* 6: 157-175.
- 5 Billich, S., Wissel, T., Kratzin, H., Hahn U., Hagenhoff, B., Lezius, A.G. and Spener F. (1988) Cloning of a full length complementary DNA for fatty acid-binding protein from bovine heart. *Eur. J. Biochem.* 175: 549-556
- 10 Binas, B., Spitzer, E., Zschiesche, W., Erdmann, B., Kurtz, A., Mueller, T., Niemann, C., Blenau, W. and Grosse, R. (1992) Hormonal induction of functional differentiation and mammary derived growth inhibitor expression in cultured mouse mammary gland explants. *In vitro Cell Dev. Biol.* 28A:625-634.
- 15 Binas, B., Gusterson, B., Wallace, R., and Clarck A.J. (1995) Epithelial proliferation and differentiation in the mammary gland do not correlate with cFABP gene expression during early pregnancy. *Dev. Gen.* 17:167-175.
- 20 Burton, P.B.J., Hogben, C.E., Jouannou, C.L., CLarck, A.G.B., Hsuan, J.J., Totty, N.F., Sorensen, C., Evans, R.W. and Tynan, M.J. (1994) Heart fatty acid binding protein is a novel regulator of cardiac myocyte hypertrophy. *Biochem. Biophys. Res. Comm.* vol 205 no 3:1822-1828.
- 25 Chevalet, C. and Corpet, F. (1986) Statistical decision rules concerning synteny or independence between markers. *Cytogenet. Cell Genet.* 43:132-139.
- Claffey, K.P., Herrera, V.L., Brecher, P. and Ruiz-Opazo, N. (1987) Cloning and tissue distribution of rat heart fatty acid-binding protein mRNA: Identical forms in heart and skeletal muscle. *Biochemistry* 26:7900-7904.
- 30 Groeneveld E (1990) PEST Users' Manual. Report-Department of Animal Sciences, University of Illinois, Urbana USA, 72 p.
- Janss, L.L.G., Van Arendonk J.A.M., Brascamp, E.W. (1994) Identification of a single gene affecting intramuscular fat in Meishan crossbreds using Gibbs sampling. *World Conference on genetics applied to livestock production*, Guelph.
- 35

- Hovenier, R., Kanis, E., Van Asseldonk, Th. and Westerink, N.G. (1992) Genetic parameters of pig meat quality traits in a halothane negative population. *Livest. Prod. Sci.* 32:309-321.
- 5 Kerr R.J. and Kinghorn B.P. (1996). An efficient algorithm for segregation analysis in large populations. *J. Anim. Breed. and Genet.* 113:457-469
- Peeters, R.A., Veerkamp, J.H., Geurts van Kessel, A., Kanda, T. and Ono, T. (1991) Cloning of the cDNA encoding human
- 10 skeletal muscle fatty acid binding protein, its peptide sequence and chromosomal localization. *Biochem. J.* 276:203-207.
- Randle, P.J., Hales, C.N., Garland, P.B., and Newsholme, E.A. (1963) The glucose fatty acid cycle: its role in
- 15 insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet* i:785-789.
- Rettenberger, G., Fries, R., Engel, W., Scheit, K.H., Dolf, G. and Hameister, H. (1994a) Establishment of a partially
- informative porcine somatic cell hybrid panel and assignment
- 20 of the loci for transition protein 2 (TNP2) and protamine 1 (PRM1) to chromosome 3 and polyubiquitin (UBC) to chromosome 14. *Genomics* 21: 558-566.
- Rettenberger, G., burkhardt, e., Adham, I.M., Engel, W., Fries, R., Klett, C. and Hameister, H. (1994b) Assignment of
- 25 the Leydig-insulin like hormone to porcine chromosome 2q12-q13 by somatic cell hybrid analysis and fluorescence in situ hybridisation. *Mamm. Genome* 5:307-309.
- Rettenberger, G., Fredholm M. and Fries, R. (1994c) Chromosomal assignment of porcine microsatellites by use of
- 30 a somatic cell hybrid mapping panel. *Anim. Genet.* 25:343-345.
- Rettenberger, G., Bruch, J., Beattie, C.W., Moran, C., Fries, R. and Hameister H. (1995a) Chromosomal assignment of
- 17 porcine microsatellites and genes by use of a somatic cell
- 35 hybrid mapping panel. *Anim. Genet.* 26: 269-273.
- Rettenberger, G., Klett, C., Zechner, U., Kunz, J., Vogel, W. and Hameister H. (1995b) Visualization of the

- conservation between pigs and humans by heterologous chromosomal painting. *Genomics* 26:372-378.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Treuner, M., Kozak, C.A., Gallahan, D., Grosse, R. and Muller, T. (1994) Cloning and characterization of the mouse gene encoding mammary-derived growth inhibitor/ heart fatty acid-binding protein. *Gene* 147:237-242.
- 10 Veerkamp, J.H., Van Kuppevelt, T.H.M.S.M., Maatman, R.G.H.J. and Prinsen C.F.M. (1993) Structural and functional aspects of cytosolic fatty acid binding proteins. *Prostaglandins Leukot. Essent. Fatty Acids* 49:887-906
- 15 Veerkamp, J.H. and Maatman, R.G.H.J. (1995) Cytoplasmic fatty acid binding proteins: their structure and genes. *Prog. Lipid Res.* vol 34 no 1:17-52.
- Wood, J.D., Enser, M., Moncrieff, C.B. and Kempster, A.J. (1988) Effects of carcass fatness and sex on the composition and quality of pigmeat. 34<sup>th</sup> International Congress of Meat Science and Technology. august 29- september 2, 1988
- 20 Brisbane Australia. 562-564.
- Yang, Y., Spitzer, E., Kenny, N., Zschiesche, W., Li, M., Kromminga, A., Muller, T., Spener, F., Lezius, A., Veerkamp, J.H., Smith, G.H., Salomon, D.S. and Grosse, R. (1994)
- 25 Members of the fatty acid-binding protein family are differentiation factors for the mammary gland. *J. Cell Biol.* vol 127 no 4:1097-1109.
- Yerle, M., Lahib-Mansais, Y., Mellink, C., Goureau, A., et al. (1995) The PIGMaP consortium cytogenetic map of the
- 30 domestic pig (*Sus scrofa domestica*). *Mamm. Genome* 9:176-186.
- Zijlstra, C., Bosma, A.A. and de Haan, N.A. (1994a) Comparative study of pig-rodent somatic cell hybrids. *Anim. Genet.* 25: 319-327.
- Zijlstra, C., Bosma, A.A., de Haan, N.A., Mellink, C.H.M. and
- 35 Groenen, M.A.M. (1994b) Confirmation of the chromosomal assignment of eight polymorphic microsatellites in the pig

using a panel of cytogenetically characterized pig-rodent somatic cell hybrids. Anim. Genet. 25 suppl.2:58.

Zijlstra, C., Bosma, A.A., de Haan, N.A. and Mellink, C.H.M. (1994c) Cytogenetic and biochemical characteristics of a

5 partly informative panel of pig-rodent somatic cell hybrids. Proc. 11th Europ. Coll. Cytogenet. Domest. Anim., Copenhagen, 170-173.

- Zijlstra, C., Bosma, A.A., de Haan, N.A. and Mellink, C.H.M. (in prep.) Construction of a cytogenetically characterized  
10 porcine somatic cell hybrid panel and its use as a mapping tool.



Table 1: Primer sequences and the corresponding size and annealing temperatures for PCR reactions of different regions of the porcine H-FABP gene.

PCR	position	primer sequence	T <sub>a</sub> (°C)	size (bp)
PCR#1	exon 1	5'GCCAGCATCACTATGGTGGACGCTTTC	57	4000
	exon 2	5'CTTAAAGCTGATCTCTGTGTTC		
PCR#2	exon 2	5'GGAGGCAAACTTGTTTACCTGC	57	1600
	exon 4	5'TCCTTCTCGTAAGTGCAGTGC		
PCR#3	exon 3	5'GGAGGCAAACTTGTTTACCTGC	62	1500
	intron 3	5'GTACTGGGAGCACTCTTCACTC		

Table 2: The primer sequences and combinations used for PCR-RFLP detection with the corresponding annealing temperature.

RFLP	Primer sequence	T (°C)	size (bp)
HaeIII- I/MspI	5' ATTGCTTCGGTGTGTTTGAG	57	950
	5' TCAGGAATGGGAGTTATTGG		
HinfI	5' GGACCCCAAGATGCTACCCCG	57	700
	5' CTGCATCTTTGACCAAGAGG		

Table 3: Allele frequencies of the porcine H-FABP gene *HaeIII*, *MspI* and the *HinfI* RFLPs in unrelated animals of different pig breeds.

		DL	Du	GY	Hs	Me	Pi	WP
		20	10	34	6	11	5	5
<i>MspI</i>	AA	0.95	0.20	0.65	1.00	1.00	0.80	0.40
	Aa	0.05	0.40	0.32	0.00	0.00	0.20	0.60
	aa	0.00	0.40	0.03	0.00	0.00	0.20	0.00
<i>HaeIII</i>	DD	0.05	0.20	0.06	1.00	1.00	0.40	0.00
	Dd	0.55	0.40	0.50	0.00	0.00	0.20	0.20
	dd	0.40	0.40	0.44	0.00	0.00	0.40	0.80
<i>HinfI</i>	HH	0.45	0.60	0.94	0.17	0.18	0.60	0.80
	Hh	0.50	0.20	0.06	0.33	0.55	0.20	0.20
	hh	0.05	0.20	0.00	0.50	0.27	0.20	0.00

Abbreviations for the various pigbreeds are DL: dutch Landrace; DU: Duroc; GY: Great Yorkshire; HS: Hampshire; ME: Meishan; PI: Pietrain; WP: Wild Pig. The second line indicates the number of unrelated animals tested.

Table 4: Chromosomal localization of the porcine hFABP gene by two independent cell hybrid panels.

The percentage concordance was determined by the total of equal absence/presence in both data sets divided by 20. The  $\phi$  value represents the correlation between both datasets. A gene can be assigned to a chromosome when the concordance is high and the  $\phi$  is more than 0.74 (syntenic). Values for  $\phi$  between 0.59 and 0.74 give no validation of the assignment. The rest is asyntenic. Using these conditions the probability of the assignment is 97.5% correct.

CHR.NR	locus	PANEL A		PANEL B	
		%CON	$\phi$	%CON	$\phi$
1	IFNA	62	0.16	46	-0.10
2	S0091	81	0.59	71	0.39
3	APOB	67	0.39	66	0.45
4	S0001	48	-0.03	66	0.34
5	S0092	71	0.45	50	0.06
6	RYS1	95	0.89	92	0.83
7	TNFB	62	0.33	62	0.10
8	ALB	48	0.31	62	0.36
9	S0095	57	0.40	50	0.15
10	S0038	57	0.28	50	0.03
11	CGT9	48	-0.03	58	0.17
12	GH	57	0.40	50	-0.03
13	S0076	48	-0.03	67	0.29
14	DAO	62	0.16	71	0.45
15	S0088	52	0.03	75	0.53
16	CGT6	57	0.28	58	0.10
17	ENDO	67	0.30	50	-0.15
18	CGT12	57	0.00	58	0.08
X	S0022	52	0.12	75	0.54
Y	SRY	76	0.39	58	0.00

Panel A: Rettenberger et al., 1994a

Panel B: Zijlstra et al., 1994a

Table 5 Number of animals (N), their mean and standard deviations (std) for intramuscular fat (IMF) and bodyweight (BW) difference from 110 kg at 180 days) for different fixed effects (effect).

5

	Effect	N	IMF(%)	(std)	BW	(kg) (std)
10	test farm					
	Someren	24	2.77	(1.66)	-4.87	(8.35)
	Herpen	18	2.00	(0.79)	-6.73	(7.85)
15	Slaughter week					
	1	5	2.02	(0.54)	-4.76	(8.34)
	2	19	2.25	(0.77)	-5.83	(6.81)
	4	6	2.82	(0.70)	-6.12	(8.80)
	5	5	2.20	(1.11)	-8.65	(11.61)
	6	7	3.09	(3.02)	-3.34	(9.63)
20	Slaughter month					
	jan	35	2.31	(0.79)	-6.13	(7.83)
	feb	7	3.09	(3.02)	-3.34	(9.63)
25	Sexe					
	boar	24	2.02	(0.71)	-3.44	(7.90)
	gilt	18	2.99	(1.85)	-8.63	(7.57)
30	Polymorphisms					
	MspI					
	AA	5	2.58	(0.69)	-10.27	(6.78)
	Aa	24	2.54	(1.73)	-7.23	(7.47)
	aa	13	2.18	(0.80)	-1.79	(8.01)
35	HaeIII					
	DD	5	2.58	(0.69)	-10.27	(6.78)
	Dd	22	2.57	(1.80)	-7.26	(7.53)
	dd	15	2.19	(0.76)	-1.01	(8.12)
40	HinfI					
	HH	21	2.02	(0.72)	-4.04	(9.06)
	Hh	15	3.00	(2.04)	-6.50	(7.00)
	hh	5	2.58	(0.69)	-10.27	(6.78)

Table 6 Values of significance for different fixed effects influencing intramuscular fat (IMF) and bodyweight (BW) using a large model (lrg) and using separate models per fixed effect (ind) (correcting for fixed effects test farm and sexe).

Effect	IMF (Pr > F)		BW (Pr >F)	
	lrg	ind	lrg	ind
regio	0.07	-	0.99	-
sexe	0.02	-	0.03	-
15 MspI	0.52	0.38	0.39	0.03
HaeIII	0.60	0.50	0.56	0.04
HinfI	0.55	0.43	0.59	0.30
Combined	-	0.80	-	0.11

Table 7 Predicted values for intramuscular fat (IMF) and bodyweight (BW) for the different fixed effects (effect) and their standard errors (s.e.).

5	<hr/>		
	Effect	IMF s.e.	BW s.e.
<hr/>			
10	sexe		
	boar	-0.61±0.18	4.78±2.52
	test farm		
	Someren	0.48±0.24	2.56±3.05
15	MspI*		
	AA	-0.02±0.32	-3.78±4.39
	Aa	0.00±0.00	0.00±0.00
	aa	-0.22±0.23	5.34±3.07
20	HaeIII*		
	DD	0.02±0.32	-4.32±4.36
	Dd	0.00±0.00	0.00±0.00
	dd	-0.12±0.22	4.35±2.87
25	HinfI*		
	HH	-0.30±0.21	2.16±2.84
	Hh	0.00±0.00	0.00±0.00
	hh	-0.08±0.32	-5.16±4.36

30 \* the different polymorphisms are analysed using one polymorphism in each model (the fixed effects test farm and sexe are included in all models).

Table 8 Differences (diff) between genotypes and their 90% (90) and 95% (95) confidence limits for intramuscular fat (IMF) and bodyweight (BW).

5	genotype	IMF		BW	
		diff	90	diff	95
10	aa-AA	-0.20	$\pm 0.39$	9.11	$\pm 8.23$
	dd-DD	-0.14	$\pm 0.39$	8.56	$\pm 6.48$
	HH-hh	-0.22	$\pm 0.38$	7.32	$\pm 6.43$



Table 9 Effect of different genotypes and the difference between the homozygotes in IMF% for each RFLP

	Genotype	effect	difference	P=0
5	<i>MspI</i>			
	AA	-0.373		
	Aa	-0.250	-0.373 $\pm$ 0.195	0.06
	aa	0.000		
10	<i>HaeIII</i>			
	DD	-0.379		
	Dd	-0.274	-0.379 $\pm$ 0.191	0.05
	dd	0.000		
	<i>HinfI</i>			
15	HH	0.402		
	Hh	0.048	0.402 $\pm$ 0.191	0.04
	hh	0.000		

Table 10                      Effect of different genotypes on backfat thickness (difference from population mean at 110 kg in mm)

5	<hr/>			
	Genotype	effect	difference	P=0
10	MspI			
	AA	-0.559		
	Aa	-0.346	-0.559 ± 0.209	0.01
	aa	0.000		
15	HaeIII			
	DD	-0.598		
	Dd	-0.342	-0.598 ± 0.207	0.004
	dd	-0.000		
15	HinfI			
	HH	0.380		
	Hh	0.092	0.380 ± 0.213	0.07
	hh	0.000		
<hr/>				

Table 11

Effect of different genotypes on bodyweight  
(difference from population mean at 180  
days in kg )

5				
	Genotype	effect	difference	P=0
	<i>MspI</i>			
	AA	-2.311		
10	Aa	-2.229	-2.311 ± 1.275	0.07
	aa	0.000		
	<i>HaeIII</i>			
	DD	-2.691		
	Dd	-2.300	-2.691 ± 1.258	0.03
15	dd	0.000		
	<i>HinfI</i>			
	HH	0.404		
	Hh	-0.621	0.404 ± 1.298	0.76
	hh	0.000		
20				

Table 12. Effect of different genotypes and the difference between the homozygotes in IMF% for each RFLP when corrected for backfat thickness.

5	<hr/>		
	Genotype	Effect	Difference
10	<i>MspI</i>		
	AA	-0.226	-0.226 $\pm$ 0.188
	Aa	-0.166	
	aa	0.000	
	<i>HaeIII</i>		
15	DD	-0.234	-0.234 $\pm$ 0.184
	Dd	-0.195	
	dd	0.000	
	<i>HinfI</i>		
	HH	0.316	0.316 $\pm$ 0.182
	Hh	0.006	
	hh	0.000	
<hr/>			

Table 13 Effect of different genotypes and the difference between the homozygotes in IMF% for each RFLP when corrected for backfat thickness and growth

---

5	Genotype	Effect	Difference
	<i>MspI</i>		
	AA	-0.245	-0.245 ± 0.188
	Aa	-0.192	
10	aa	0.000	
	<i>HaeIII</i>		
	DD	-0.250	-0.250 ± 0.184
	Dd	-0.214	
	dd	0.000	
15	<i>HinfI</i>		
	HH	0.316	0.316 ± 0.182
	Hh	0.001	
	hh	0.000	

CLAIMS

1. An isolated or recombinant pig H-FABP gene specific nucleic acid molecule or pig H-FABP gene specific fragments thereof comprising or hybridising to the nucleotide sequence as listed in figure 1 or its complementary sequence or the  
5 RNA equivalents thereof .
2. A method localising, identifying or marking genes or alleles or quantitative trait loci of pigs, using a molecule or a fragment or fragments thereof according to claim 1.
3. A method according to claim 2 whereby genes or alleles  
10 or quantitative trait loci are localised, identified or marked that are associated with production traits of pigs.
4. A method according to claim 2, or 3 to identify or mark alleles of the pig H-FABP gene.
5. A method according to claim 4 distinguishing between  
15 alleles of the H-FABP gene of pigs.
6. A method according to claim 5 by detecting specific restriction sites in an allele of the H-FABP gene of pigs.
7. A method according to claim 6 whereby an *MspI* restriction site is detected.
- 20 8. A method according to claim 6 whereby an *HaeIII* restriction site is detected.
9. A method according to claim 6 whereby an *HinfI* restriction site is detected.
10. A method using a molecule, a fragment or fragments  
25 thereof according to claim 1, localising, identifying or marking genes or alleles or quantitative trait loci in samples, in particular biological samples, cells or tissues, such as but not limited to hair, skin or blood, by allowing for specific amplification of genomic fragments of those  
30 genes or alleles or quantitative trait loci.
11. A diagnostic assay or kit according to claim 10.
12. A method according to anyone of claims 2-9 localising, identifying or marking genes or alleles or quantitative trait loci in samples, in particular biological samples,

cells or tissues, such as but not limited to hair, skin or blood of pigs, by allowing for specific amplification of genomic fragments of those genes or alleles or quantitative trait loci.

- 5 13. A method according to anyone of claims 10 or 12 in which the method of amplification is the polymerase chain reaction.
14. Diagnostic assay or kit according to claim 13.
15. A method according to anyone of claims 2-14 identifying  
10 differences between alleles of the pig that are associated with differences in production traits of pigs.
16. A method according to claim 15 identifying alleles of the pig that are associated with improved production traits of the pig.
- 15 17. Use of the methods according to anyone of claims 2-16 in marker assisted identification of pigs or in marker assisted selection of pigs.
18. Use of the methods according to claims 16 or 17 in breeding programmes.

aagctttttaaattctttgagctatttaggggattactgaaactttccatatgggggagagaagaacc  
 atcagccccactgtccttggagccttcagaaacacacaggaatttcctacccccatgaccagg  
 gtaatatattacttaagaggctgcaagcattggtaaagtggctcaagatttggagtcagaact  
 cagggtaaataaattctgactttcctttgttgtagttaaagtcattttaaacaagttgttttat  
 ttgccttggagcctctctttcctcagctctaaaaatggggaggacccaaaaggccatgcagttg  
 ggagcctgagaaaatcagatgtggaatccctgcatagagcctggtaacgggcttacatccacc  
 cttgggatgtactctgcttccctccttgaaggcagtgaccaagcagcatgtccagcttcccc  
 tgaggcagcatctctgttttctcagaatgtgaagagtacaaagacctcattgaatccgtgctg  
 gaggaggaagtgtgttttcaggatggagagctggcagagaagatgagcctggctgcaaggctt  
 gggtaaggagggtacctgtgtgggaaagtatgtgaagtggggaagcccaagctggaagaataaa  
 gagataatttgtaccaggagagaagcagaaaatgtagccggtaggcacagaattagacaaaaga  
 cagcttgtgggcctgtgggaactggatgggggcctagttagccctctagattcacccttaaat  
 aatgcatgaataaaaaggagacagacgcctgggttaaaaagactagcaacatggcttctagctt  
 tgtgacctggggcaaatttacttaactttcctcagttacctgagccacctcatttgaagatg  
 aaaataccaacagctctaccttagaggggttatgaagattaaatgtgattacaagtagttagaag  
 catctcatatcatgcatgaacaaatatttgcgtgaattattatacttgttcttagtttttatag  
 gtttggtaattatcgcccttttaaaaaatgttagatctggactctacctcaccctctccttaaat  
 cctgcgaccttgtcagtcacctctaaagtcttggactcctcacctgtaaaaagaggaccaagat  
 gcctacgcgcgagagagtcggccagaactttcccagcggggcatcagcccaagagtgaagttc  
 ctttctcctctagctctctcatctctgttcggacccagggctcgtggagggaaccaggaaaccga  
 agatgtccatgacctgcgccccaaactcgcgggctccgtgccccagcggcttccctctcagat  
tccgaagagcgcccttgaggccaggaaaggggactggctcctgccagccccggggagggttcgg  
 atccccgggtacccgtgctgcccggcgccctacgggctgcccccttctctcttcggcaggctggg  
 gatgggcatcaaccggcgaggcagcgccctgcatggggcttccctatttccggagcggggcg  
 tacgccacgcctcgtcacgtgacgctagggccattttaaagcggtagcgcgggctgggagccgc  
 cggctcctggaatttttgcgcgcctgttctgtcgtctctttctcagcctagcccagcctcaccA  
**TGGTGGACGCCTTCGCGGGCACCTGGAAGCTAGTGGACAGCAAGAATTTGATGACTACATGA**  
**M V D A F A G T W K L V D S K N F D D Y M**

**AGTCAATTG**gtgaggacgactcgggatgttgggctgggaagaggttggccgcgtgccccggccg  
**K S I**

gagccttgcctccagccctggctgggaccctgtgnnn<3.5kb>ctctgccattgaacattctg  
 ggcttgaattccagatttggactctgggcttcaacttactctgtttttgcatctgtagagtta  
 ataacacagatccttcaggagactggaagaattgaatgagattacatgtgtagttgttctac  
 tacacagaaagcacttaatagttttttttttttttttttttttttttttttgtcttttttagggccata  
 tccatggcacatggagttcccaagctaggggtccaatcagagctgaagcgctagcttttgcca  
 gagccacagcagcaggggttccgagccctgtccttaaccacccaagcgaagccagggttgaa  
 cccgcctcctcatggatactagttgagtttgttaaccactgagccacgatgttaactcccaat  
 ggtagcttattctaatctcggctctccactgacttgcgtgtgargcatgcctgttcccttaagc  
 ttctcatttgtctaatgggatcagtaatgtctcttcatagcacagagctacgaagactgaatt  
 taaaaacagccatgcagcctttgaaaattcttgcctcctccttcttcttccacatctagg  
 cctaagagctgtttgtgtctgcgggtctgttccagcttgtactcataaccttccctctgcc  
 cttag**GTGTGGGTTTTGCCACCAGGCAGGTGGCCAACATGACCAAGCCTACCACAATCATCGA**  
**G V G F A T R Q V A N M T K P T T I I E**

**AGTGAATGGGGACACAATCATCATAAAAACACAAAGCACCTTCAAGAGCACAGAGATCAGCTT**  
**V N G D T I I I K T Q S T F K S T E I S F**

**CAAGCTGGGAGTGGAGTTTGATGAGACAACAGCAGATGACAGGAAGGTCAAG**gtgagttgagg  
**K L G V E F D E T T A D D R K V K**

gaagtgtcacagggaatgaaagtccttgaagggggaataggatggcaccctattattgcacctt  
 gaggggtaggatgttatgaggaattgaaacaaagtccttaagcctgtgcagtgctgggatgt

Figure 1/1

**UBSTITUTE SHEET (RULE 2 )**



2/2

cctgagtgctaaaagcatgtcctgtcctcttgttttctcctccctcccttcttgggaagcatcta  
 cctgttacctctggacgagggcagcatcaagccaggatattctgacctccgtgtctgctgcag  
 ctccagctgggtgaccttgtgcctggcatgcaacacctctggctttcttcccccttccataga  
 tgggggttatgtggacatggctgtaacctcaagtgttttaggacaatgccctgaagtcaaatac  
 ctctgctaaataggccatcctggcctctagcagcagagacagctgggggcaagagggtgaaac  
 ctactcagatgcttctgcttctccagccaccacctcttgggcttggccagtgacctgaggaa  
 ctgtttccaaggctgccagcccacaagcaggdaaaactatgttctggatggagcattgcttcg  
 gtgtgtttgagtgcttttctgccaaagcctcagctccccctccttccaaggaggaggggacatc  
 taccctctctcaggatcggctaccctgcagcagcttgccctcaggcctcccagccccctgggcca  
 ggcattggaagaggagaaactgggagggttctctctacagggtgtccctatagtagtatcttt  
 tcaccagaccgatcattcagctactcagctgtttcctcttccctcacagctgcctccaatagt  
 ttccctccttgtggaggaagcgggcagcagagatgggacagtagccagagatggagagggtgct  
 gtccctaccacatgcttttaacttaaaagttagcctaccttcttgaggggcaagaaatccaattaa  
 acgccaacagttctatgggatggccagggtagaaataatctgctctgagtggaacaaaagctg  
 ggggaatgggaatgaggaagggtgatcataagggtgtacccctagccctaaacacacacagcct  
 tttcagattctcctattaagcctgctcagtcacgggtcccctagcctggcttagagccccaga  
 cttctagaagagacacctgggagagctgccaaatgagaaccagctgatgtggactggcagcag  
 caccagaagagtttaagtggacatttaaccagggtgtggaacactccaagggtacccagcc  
 ccaggacaaggcagttgctgccacttgtcagctgaagggtcacatgatatggaccaagagg  
 agtaggaatacagggcagagtcctaattttccaataactcccattcctga<0.4kb>gagctc  
 tggctcctagcaccatcaacagtttaggttagaggctgagctctgctgtggccagagtgaccaga  
 cacccttctctgtccctcaatcccagctcaggcctctaccctctttccacagTCCATTGTGA  
 S I V

CACTGGATGGAGGCAAACCTTGTCACCTGCAGAAGTGGAATGGACAAGAGACAACGCTTGTTCT  
 T L D G G K L V H L Q K W N G Q E T T L V

GGGAAGTAGTTGATGGGAAACTCATCTGgttaagatggacaactttggagttcaccagttg  
 R E L V D G K L I L

gttcttactgccccctcagccaaccctactgtgaaaagccaagggttccctgggaccagtgga  
 gctgggtagtgagggtggaggcaaggctgggtggcattaggttaggaattttctt  
 atgtagtggctttggactcacacaagattgggttttaatacacagtttataggctacatgactt  
 ggggtaagttaggttctccatgtccaactgtaaagtgggtattaatgacaccagccttggagg  
 gctgctgtgaagtctgagggagatgggtggatgaggagcacaggat<1.0kb>tacgagaggaa  
 ttacagtgtgaacttgaagacctgggtgaagcattaggaaaaatacatgtcaagtacctgg  
 catgtagtaggtagtaaatgacgatattagcatgaaggtagagggtttccccggcagagtga  
 agagtgtcccagtagccttctgtgcccttaccctgccctaattctgacctcttttattccag  
 AACTCACCCATGGCAGTGCAGTTTGCACCTCGCACTTACGAGAAAGAGGCATGAacctgccat  
 T L T H G S A V C T R T Y E K E A stop

cccttcgactgttccctgccaattggctactcctggactcagcaccagattgcctcatttttct  
 ctctggcattttgtaaaaatctactttggggatattctcctggggtcagggttgaccagcctg  
 cgttcagttccggttcttgtgtgtatgttagttttttttaattgcatccaaagggtgctct  
 gaggtcaataaaatagccaaggc

Figure 1/2





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 14/47, C12Q 1/68, C07K 14/705</b>		<b>A3</b>	(11) International Publication Number: <b>WO 97/35878</b>
			(43) International Publication Date: 2 October 1997 (02.10.97)
(21) International Application Number: PCT/NL97/00157 (22) International Filing Date: 27 March 1997 (27.03.97) (30) Priority Data: - 96200855.3 28 March 1996 (28.03.96) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicants (for all designated States except US): DALLAND B.V. [NL/NL]; Postbus 128, NL-5740 AC Beek en Donk (NL). PROVA B.V. [NL/NL]; Postbus 290, NL-5280 AG Boxtel (NL). STAMBOEK ZUID B.V. [NL/NL]; Postbus 86, NL-5268 ZH Helvoirt (NL). NOORD NEDERLANDS VARKENSSTAMBOEK B.V. [NL/NL]; Hoofdweg 2, NL-7382 BH Klarenbeek (NL). INSTITUUT VOOR DIERHOUDERIJ EN DIERGEZONDHEID (ID-DLO) [NL/NL]; Edelhertweg 15, NL-8219 PH Lelystad (NL). (72) Inventor; and (75) Inventor/Applicant (for US only): GERBENS, Frans [NL/NL]; Botter 25-02, NL-8232 KP Lelystad (NL). (74) Agent: SMULDERS, Th., A., H., J.; Vereenigde Octrooibureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. (88) Date of publication of the international search report: 30 October 1997 (30.10.97)	
(54) Title: THE PORCINE HEART FATTY ACID-BINDING PROTEIN ENCODING GENE AND METHODS TO IDENTIFY POLYMORPHISMS ASSOCIATED WITH BODY WEIGHT (57) Abstract <p>The present invention provides a novel sequence of the pig H-FABP gene, as well as methods of using said gene and its products. Especially the invention provides methods for detecting different alleles of the pig H-FABP gene, which different alleles are associated with differences in the genotypic and/or phenotypic traits of the pigs having those alleles. Especially the invention provides methods for distinguishing between alleles resulting in different phenotypes, particularly using techniques involving selective amplification of pig H-FABP gene derived materials. These techniques are especially suitable for selecting animals to be used in breeding programmes. Breeding programmes employing such techniques are also disclosed.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/NL 97/00157

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07K14/47 C12Q1/68 C07K14/705

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07K C12Q A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BIOCHEMICAL JOURNAL, vol. 276, 1991, LONDON GB, pages 203-207, XP000196430 R.A. PEETERS ET AL.: "Cloning of the cDNA encoding human skeletal-muscle fatty-acid binding protein, its peptide sequence and chromosomal localization" cited in the application see the whole document --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*I\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*A\* document member of the same patent family

Date of the actual completion of the international search

2 July 1997

Date of mailing of the international search report

16.09.1997

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340 3016

Authorized officer

De Kok, A

## INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/NL 97/00157

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, vol. 264, no. 2, 1 August 1988, SAN DIEGO US, pages 533-545, XP000196455 R.J. PAULUSSEN ET AL.: "Characterization and binding properties of fatty-acid binding proteins from human, pig, and rat heart" see the whole document ---	1
A	DE 43 38 817 A (MAX DELBRÜCK CENTRUM FÜR MOLEKULARE MEDIZIN) 14 June 1995 see the whole document ---	1
A	WO 92 18651 A (IOWA STATE UNIVERSITY RESEARCH FOUNDATION, INC.) 29 October 1992 see the whole document ---	1-11
A	ANIMAL GENETICS, vol. 22, no. 5, 1991, OXFORD GB, page 443 XP000196444 W BARENDSE ET AL.: "TaqI reveals a polymorphism in cattle when probing with the rat fatty acid binding protein (I-FABP)" see the whole document ---	1,4-6
A	BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1081, no. 1, 1991, AMSTERDAM NL, pages 1-24, XP000196446 J.H. VEERKAMP ET AL.: "Structural and functional features of different types of cytoplasmic fatty acid-binding proteins" see the whole document -----	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 97/00157

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 4338817 A	14-06-95	NONE	
WO 9218651 A	29-10-92	AU 660994 B	13-07-95
		AU 1917692 A	17-11-92
		BR 9205918 A	08-11-94
		CA 2108579 A	20-10-92
		EP 0580767 A	02-02-94
		JP 6506837 T	04-08-94
		US 5550024 A	27-08-96
		US 5374526 A	20-12-94

